

Noncommutative quantum mechanics of a test particle under linearized gravitational waves

Anirban Saha*

*Department of Physics and Astrophysics,
West Bengal State University,
Barasat, North 24 Paraganas,
West Bengal, India*

and

*Visiting Associate in Inter University Centre
for Astronomy and Astrophysics, Pune, India*

Sunandan Gangopadhyay†

*Department of Physics and Astrophysics,
West Bengal State University,
Barasat, North 24 Paraganas,
West Bengal, India*

and

*Visiting Associate in S.N.Bose National Centre
for Basic Sciences, Kolkata, India*

(Dated: October 8, 2009)

We consider the quantum dynamics of a test particle in noncommutative space under the influence of linearized gravitational waves in the long wave-length and low-velocity limit. A prescription for quantizing the classical Hamiltonian for the interaction of gravitational wave with matter in noncommutative space is proposed. The Hamiltonian (and hence the system) is then exactly solved by using standard algebraic methods. The solutions show prominent signatures of the noncommutative nature of space. Computation of the expectation value of the particle's position reveals the inherent quantum nature of spacetime noncommutativity.

PACS numbers: 11.10.Nx, 03.65.Ta, 11.10.Ef, 04.30.Nk, 42.50.Dv

Gravitational waves (GWs) are tiny disturbances in spacetime. Their effect on matter is expected to be measured by the relative optical phase shift between the light paths in two perpendicular km-length arm cavities caused by the tiny displacement of two test mass mirrors (hung at the end of each cavity) induced by passing GWs. The typical amplitude of GWs $h \sim \frac{\delta L}{L}$ emitted by binary systems in the VIRGO cluster of galaxies, at a distance $\sim 20\text{Mpc}$, at 100Hz which the ground based Laser-interferometric GW detectors like LIGO¹, VIRGO², GEO³ TAMA⁴ etc. are designed to probe is $\sim 10^{-21}$. Since the cavity arm length $L \sim 1\text{km}$, δL will be $\sim 10^{-18}\text{m}$. It is, therefore, at the quantum mechanical level that experimental evidence for the GWs is likely to appear⁵.

Interestingly, in recent developments of noncommutative (NC) quantum mechanics and NC quantum field theory, where the coordinates x^μ satisfy the NC algebra

$$[x^\mu, x^\nu] = i\theta^{\mu\nu} \quad (1)$$

the upperbounds on various NC parameters appearing in the literature^{7,8,9,10,11,12,25} are quite close to this length scale. A wide range of theories have been constructed in a NC framework including various gauge theories¹³, gravity¹⁴ and even encompassing certain possible phenomenological consequences^{16,17,18,19,20}. The upperbound on the value of the coordinate commutator θ^{ij} found in²⁵ is $\lesssim (10\text{TeV})^{-2}$ which corresponds to

$4 \times 10^{-40}\text{m}^2$ for $\hbar=c=1$. Whereas such upperbounds on time-space NC parameter θ^{0i} is $\lesssim 9.51 \times 10^{-18}\text{m}^2$. However, recent studies in NC quantum mechanics revealed that the NC parameter associated with different particles are not same²¹ and this bound could be as high as $\theta \lesssim (4\text{GeV})^{-2} - (30\text{MeV})^{-2}$ ¹⁰. These upperbounds correspond to the length scale $\sim 10^{-20}\text{m} - 10^{-17}\text{m}$.

With the prospect of the direct detection of GWs of such tiny amplitude as $\sim 10^{-18}$ in the near future, a sharp possibility of detecting the NC structure of spacetime would be in the GW detection experiments. Therefore, analyzing the interplay of classical GWs with a test particle in a *NC quantum mechanical* framework becomes important in its own right as it may predict some detectable signature of noncommutativity.

In light of these observations, we would like to construct the quantum mechanics of a test particle in NC spacetime, interacting with a linearised GW in the long wave-length and low velocity limit. Since it has been demonstrated in various formulations of NC general relativity^{14,15} that any NC correction in the gravity sector is second order in the NC parameter, therefore, in a first order theory in NC space, the linearised GW remains unaltered by NC effects and any NC correction appearing in the system will be through the particle sector only. This is true not only with the canonical (i.e. constant) form of noncommutativity but also for the Lie-algebraic NC spacetime¹⁵. We shall incorporate the NC

effect in the problem by writing the NC Hamiltonian and then reexpressing it in terms of the commutative coordinates and their momenta by the well known Bopp-shift transformations^{10,12}. Then we shall quantize the system following²². As a first step towards the formulation of a NC quantum mechanics of GW interacting with a test particle, we shall concentrate on the simplest of the GWs, namely the linearly polarized ones in this analysis.

In a linearised theory of gravity, once we choose the transverse-traceless (TT) gauge, all the gauge redundancies of the theory get removed and the GW is characterised by the only non-zero components²³ $h_{11} = -h_{22}$, and $h_{12} = h_{21}$, called the $+$ and $\times$ polarisation respectively. The only non-trivial components of the curvature tensor in TT-gauge are²⁷

$$R^j_{0,k0} = -\frac{\partial \Gamma^j_{0k}}{\partial t} = -\frac{1}{2} \frac{\partial^2 h_{jk}}{\partial t^2} \quad (2)$$

and the geodesic deviation equation in the proper detector frame becomes²³

$$m \frac{d^2 x^j}{dt^2} = -m R^j_{0,k0} x^k \quad (3)$$

which governs the response of a scalar spin-zero test particle to the passage of a GW. Here x^j is the proper distance of the test particle from the origin and m is its mass. For convenience, we have assumed that GW detectors can be reasonably isolated so that any external forces other than the GW interaction are negligible, i.e. the particles are considered to be free otherwise. Also, the GW is treated as an external classical field. Eq.(3) can be used to describe the evolution of proper distance in TT-gauge frame as long as the spacial velocities involved are non-relativistic. Also, $|x^j|$ has to be much smaller than the typical scale over which the gravitational field changes substantially, i.e. the reduced wavelength $\frac{\lambda}{2\pi}$ of GW. This situation is referred to as the *small-velocity and long wavelength limit*. Thus, with Eq.(3) we can analyze the interaction of GW with a detector²⁸ which has a characteristic linear size $L \ll \frac{\lambda}{2\pi}$.

The Lagrangian for the system, whose time evolution is described by Eq. (3), can be written, upto a total derivative term²² as

$$\mathcal{L} = \frac{1}{2} m \dot{x}^2 - m \Gamma^j_{0k} \dot{x}_j x^k. \quad (4)$$

The canonical momentum corresponding to x_j is $p_j = m \dot{x}_j - m \Gamma^j_{0k} x^k$ and the Hamiltonian becomes

$$H = \frac{1}{2m} \left(p_j + m \Gamma^j_{0k} x^k \right)^2. \quad (5)$$

If we take GWs propagating along the z -axis, then due to the transverse nature of GWs, Γ^j_{0k} has non-zero components only in the $x-y$ plane and the particle undergoes free motion along the z -direction. Hence we can essentially focus on the planar motion of the particle.

To “quantize” this system in the NC plane, we now replace x^j and p_j in the above Hamiltonian by operators $\hat{x}^j$ and $\hat{p}_j$ satisfying the NC Heisenberg algebra²⁹

$$[\hat{x}_i, \hat{p}_j] = i\hbar \delta_{ij}, \quad [\hat{x}_i, \hat{x}_j] = i\theta \epsilon_{ij}, \quad [\hat{p}_i, \hat{p}_j] = 0. \quad (6)$$

It is well known that this can be mapped to the standard ($\theta = 0$) Heisenberg algebra spanned by X_i and P_j using^{10,12}

$$\hat{x}_i = X_i - \frac{1}{2\hbar} \theta \epsilon_{ij} P_j, \quad \hat{p}_i = P_i. \quad (7)$$

Using the traceless property of the GW and rewriting the NC version of Eq.(5) in terms of the operators X_i and P_j , we obtain

$$\hat{H} = \frac{P_j^2}{2m} + \Gamma^j_{0k} X_j P_k - \frac{\theta}{2\hbar} \epsilon_{jm} P_m P_k \Gamma^j_{0k}. \quad (8)$$

Observe that a direct coupling between the GW and spatial noncommutativity appears here. Since we are dealing with linearized gravity, a term quadratic in Γ has also been neglected in Eq.(8).

It is to be noted that the above Hamiltonian can be obtained from the following Lagrangian (which can be computed from the Hamiltonian (8) by an inverse Legendre transformation)

$$\hat{\mathcal{L}} = \frac{m}{2} \left[(\dot{X}_i^2 - 2\Gamma^j_{0i} \dot{X}_i X_j) + \frac{\theta m}{\hbar} \epsilon_{jk} \Gamma^j_{0i} \dot{X}_i \dot{X}_k \right]. \quad (9)$$

This is reassuring since we do have both the Lagrangian and the Hamiltonian of the problem in hand as in the commutative ($\theta = 0$) case. Further, the above form of the Lagrangian can be obtained from the noncommutative version of the Lagrangian (4) by making the following Bopp shift

$$\hat{x}_i = X_i - \frac{1}{2\hbar} \theta \epsilon_{ij} P_j, \quad \hat{\dot{x}}_i = \dot{X}_i \quad (10)$$

and then putting $P_i = m \dot{X}_i$ since both the quadratic terms in θ and Γ can be neglected. This can be regarded as the Lagrangian version of the Bopp shift (7) applied at the Hamiltonian level.

Defining raising and lowering operators

$$X_j = \left(\frac{\hbar}{2m\varpi} \right)^{1/2} (a_j + a_j^\dagger) \quad (11)$$

$$P_j = -i \left(\frac{\hbar m \varpi}{2} \right)^{1/2} (a_j - a_j^\dagger) \quad (12)$$

where ϖ is determined from the initial uncertainty in either the position or the momentum of the particle, we write the Hamiltonian (8) as

$$\begin{aligned} \hat{H} = & \frac{\hbar \varpi}{4} \left(2a_j^\dagger a_j + 1 - a_j^2 - a_j^{\dagger 2} \right) - \frac{i\hbar}{4} \dot{h}_{jk} \left(a_j a_k - a_j^\dagger a_k^\dagger \right) \\ & + \frac{m\varpi\theta}{8} \epsilon_{jm} \dot{h}_{jk} \left(a_m a_k - a_m a_k^\dagger + C.C \right) \end{aligned} \quad (13)$$

where C.C means complex conjugate. Working in the Heisenberg representation, the time evolution of $a_j(t)$ is given by

$$\begin{aligned} \frac{da_j}{dt} = & -i\frac{\varpi}{2}(a_j - a_j^\dagger) + \frac{1}{2}\dot{h}_{jk}a_k^\dagger \\ & + \frac{im\varpi\theta}{8\hbar}(\epsilon_{lj}\dot{h}_{lk} + \epsilon_{lk}\dot{h}_{lj})(a_k - a_k^\dagger) \end{aligned} \quad (14)$$

and that of $a_j^\dagger(t)$ is the C.C of Eq.(14). Next, noting that the raising and lowering operators must satisfy the commutation relations

$$[a_j(t), a_k(t)] = 0, \quad [a_j(t), a_k^\dagger(t)] = \delta_{jk} \quad (15)$$

we write them in terms of $a_j(0)$ and $a_j^\dagger(0)$, the free operators at $t = 0$, by the time-dependent Bogoluibov transformations

$$\begin{aligned} a_j(t) &= u_{jk}a_k(0) + v_{jk}a_k^\dagger(0) \\ a_j^\dagger(t) &= a_k^\dagger(0)\bar{u}_{kj} + a_k(0)\bar{v}_{kj} \end{aligned} \quad (16)$$

where the bar denotes the C.C and u_{jk} and v_{jk} are the generalized Bogoluibov coefficients. They are 2×2 complex matrices which, due to eq. (15), must satisfy $uv^T = u^T v$, $uu^\dagger - vv^\dagger = I$, written in matrix form where T denotes transpose, $\dagger$ denotes conjugate transpose and I is the identity matrix. Since $a_j(t=0) = a_j(0)$, u_{jk} and v_{jk} have the boundary conditions $u_{jk}(0) = I$ and $v_{jk}(0) = 0$. Then, from eq.(14) and its C.C we get the equations of motions in terms of $\xi = u + v^\dagger$ and $\zeta = u - v^\dagger$:

$$\frac{d\xi_{jk}}{dt} = -i\varpi\zeta_{jk} + \frac{\dot{h}_{jl}}{2}\xi_{lk} + \Theta_{jl}\zeta_{lk} \quad (17)$$

$$\frac{d\zeta_{jk}}{dt} = -\frac{1}{2}\dot{h}_{jl}\zeta_{lk} \quad (18)$$

where Θ_{jl} is the new term reflecting the interplay of non-commutativity with GW

$$\Theta_{jl} = \frac{im\varpi\theta}{4\hbar}(\dot{h}_{jm}\epsilon_{ml} - \epsilon_{jm}\dot{h}_{ml}). \quad (19)$$

Eq(s). (17, 18) are difficult to solve analytically for general h_{jk} . However, our goal, in the present letter, is to investigate to what extent spatial noncommutativity affects the interaction of GWs with spin-zero test particle in the simplest of settings. Therefore we shall solve Eq(s). (17, 18) for the special case of linearly polarized GWs.

In the two-dimensional plane, the GW, which is a 2×2 matrix h_{jk} , is most conveniently written in terms of the Pauli spin matrices as

$$h_{jk}(t) = 2f(t)(\epsilon_\times\sigma_{jk}^1 + \epsilon_+\sigma_{jk}^3) = 2f(t)\epsilon_A\sigma_{jk}^A. \quad (20)$$

Note that the index A runs from 1–3, however, no contribution from σ^2 is included. $2f(t)$ is the amplitude of the GW whereas $\epsilon_\times(t)$ and $\epsilon_+(t)$ represent the two possible polarization states of the GW and satisfy the condition

$\epsilon_\times^2 + \epsilon_+^2 = 1$ for all t . In case of linearly polarized GWs however, the polarization states ϵ_A are independent of time and $f(t)$ is arbitrary. To set a suitable boundary condition we shall assume that the GW hits the particle at $t = 0$ so that

$$f(t) = 0, \quad \text{for } t \leq 0. \quad (21)$$

We now move on to solve Eq.(18) by noting that any 2×2 complex matrix can be written as a linear combination of the Pauli spin matrices and identity matrix. Hence we make the ansatz :

$$\zeta_{jk}(t) = A(t)I_{jk} + B(t)\epsilon_A\sigma_{jk}^A. \quad (22)$$

Substituting for h_{jk} and ζ_{jk} from Eq.(20) and Eq.(22) respectively in Eq.(18) and comparing the coefficients of I and $\epsilon_A\sigma^A$, we get first order differential equations for A and B which can be readily integrated to obtain

$$\zeta_{jk} = \cosh[f(t)]I_{jk} - \sinh[f(t)]\epsilon_A\sigma_{jk}^A \quad (23)$$

once we have used the boundary conditions Eq.(21). Now that we have the solution for ζ , we can take a closer look at the NC correction term in Eq.(19), before proceeding to solve Eq.(17). Computation shows that the coupling between noncommutativity and GW not only deforms the effects of the two polarization states $\epsilon_\times\sigma_{jk}^1$ and $\epsilon_+\sigma_{jk}^3$ present in the GW but also generates a new term proportional to the Pauli spin matrix σ_{jk}^2 :

$$\Theta_{jl}\zeta_{lk} = \frac{m\varpi\theta}{\hbar}\dot{f}(t)\left\{iA(\epsilon_+\sigma^1 - \epsilon_\times\sigma^3)_{jk} + B\sigma_{jk}^2\right\} \quad (24)$$

Hence, as a trial solution we take a more general form for ξ_{jk} that accomodates all the Pauli matrices along with the identity matrix

$$\xi_{jk}(t) = CI_{jk} + D_1\sigma_{jk}^1 + D_2\sigma_{jk}^2 + D_3\sigma_{jk}^3. \quad (25)$$

Substituting for h_{jk} , ζ_{jk} and ξ_{jk} from Eq(s).(20), (22) and (25) respectively in Eq.(17) and comparing the coefficients of I and σ^A , we get the following set of first order differential equations

$$\begin{aligned} \dot{C} &= -i\varpi A + \dot{f}(D_1\epsilon_\times + D_3\epsilon_+) \\ \dot{D}_1 &= -iB\varpi\epsilon_\times + \dot{f}\left(C\epsilon_\times - iD_2\epsilon_+ + \frac{im\varpi\theta}{\hbar}A\epsilon_+\right) \\ \dot{D}_2 &= i\dot{f}(D_1\epsilon_+ - D_3\epsilon_\times) + \frac{m\varpi\theta}{\hbar}\dot{f}B \\ \dot{D}_3 &= -iB\varpi\epsilon_+ + \dot{f}\left(C\epsilon_+ + iD_2\epsilon_\times - \frac{im\varpi\theta}{\hbar}A\epsilon_\times\right) \end{aligned} \quad (26)$$

With the recombination

$$G = D_1\epsilon_\times + D_3\epsilon_+, \quad Q = D_1\epsilon_+ - D_3\epsilon_\times \quad (27)$$

we recast the set of Eq(s).(26) in the following form

$$\begin{aligned} \dot{C} &= -i\varpi A + \dot{f}G, \quad \dot{G} = -i\varpi B + \dot{f}C \\ \dot{D}_2 &= i\dot{f}Q + \frac{m\varpi\theta}{\hbar}\dot{f}B, \quad \dot{Q} = -\dot{f}D_2 + \frac{m\varpi\theta}{\hbar}i\dot{f}A \end{aligned} \quad (28)$$

Integrating Eq(s). (28) and (29) and imposing the boundary condition Eq.(21) we get

$$\begin{aligned} C &= \cosh f(t) - \frac{i\varpi}{2} F_+(t) ; \quad G = -\frac{i\varpi}{2} F_-(t) ; \\ D_2 &= -\frac{m\varpi\theta}{\hbar} f(t) \sinh[f(t)] ; \\ Q &= -\frac{im\varpi\theta}{\hbar} f(t) \cosh[f(t)] ; \\ F_{\pm}(t) &= \left[e^{f(t)} \int_0^t dt' e^{-2f(t')} \pm e^{-f(t)} \int_0^t dt' e^{2f(t')} \right] . \end{aligned}$$

Inverting the set of equations in (27), we can express D_1 and D_3 in terms of G and Q . Combining the expressions for C , D_1 , D_2 and D_3 we can write the solution for ξ using Eq(s) (25) which along with ζ in Eq. (23), give u_l and v_l , where l stands for linear polarization :

$$u_l(t) = I \left[\cosh f - i\frac{\varpi}{4} F_+ \right] - \frac{i\varpi}{4} (\epsilon_{\times}\sigma^1 + \epsilon_{+}\sigma^3) F_- + \left(\frac{m\varpi\theta}{2\hbar} \right) f \{ i (\epsilon_{+}\sigma^1 - \epsilon_{\times}\sigma^3) \cosh f + \sigma^2 \sinh f \} , \quad (30)$$

$$v_l(t) = I \frac{i\varpi}{4} F_+ + (\epsilon_{\times}\sigma^1 + \epsilon_{+}\sigma^3) \left[\sinh f + i\frac{\varpi}{4} F_- \right] - \left(\frac{m\varpi\theta}{2\hbar} \right) f \{ i (\epsilon_{+}\sigma^1 - \epsilon_{\times}\sigma^3) \cosh f + \sigma^2 \sinh f \} . \quad (31)$$

The system has now been essentially solved for the linearly polarized GWs once one specifies the initial expectation values of the particle position and momentum and also the initial uncertainty in either the position or the momentum of the particle to determine ϖ . For example, one can assume the initial wave-function³⁰ of the particle to be a Gaussian wave-packet

$$\psi(x, y) = \frac{1}{\sqrt{\pi}l_0} e^{-\frac{1}{2l_0^2}[(x-x_0)^2 + (y-y_0)^2]} e^{\frac{i\vec{p}_0 \cdot \vec{r}}{\hbar}} \quad (32)$$

located at (x_0, y_0) and moving with momentum $\vec{p}_0 = (p_{x_0}, p_{y_0})$ when the GW just hits the system at time $t = 0$. This will give uncertainties in momentum and position as $\sqrt{\frac{\hbar m \varpi}{2}}$ and $\sqrt{\frac{\hbar}{2m\varpi}}$ respectively. Either can be used to compute ϖ . From the initial position and momentum expectation values, i.e. $\langle \vec{r}_0 \rangle = (x_0, y_0)$ and $\langle \vec{p}_0 \rangle = (p_{x_0}, p_{y_0})$, we get the raising and lowering operator at time $t = 0$. We then use Eq(s) (16), (30) and (31) to find $a_j(t)$ and $a_j^\dagger(t)$ at a general time. This in turn gives us the expectation value of the position of the particle at any arbitrary time t

$$\begin{aligned} \langle x(t) \rangle &= \cosh f(t) x_0 + (\epsilon_{+} x_0 + \epsilon_{\times} y_0) \sinh f(t) + (F_+ \\ &+ \epsilon_{+} F_-) \frac{p_{x_0}}{2m} + \epsilon_{\times} F_- \frac{p_{y_0}}{2m} + \frac{\theta p_{x_0}}{\hbar} \epsilon_{\times} f(t) \cosh f(t) \\ &+ \frac{\theta p_{y_0}}{\hbar} [f(t) \sinh f(t) - \epsilon_{+} f(t) \cosh f(t)] \quad (33) \end{aligned}$$

$$\begin{aligned} \langle y(t) \rangle &= \cosh f(t) y_0 + (\epsilon_{\times} x_0 - \epsilon_{+} y_0) \sinh f(t) + (F_+ \\ &- \epsilon_{+} F_-) \frac{p_{y_0}}{2m} + \epsilon_{\times} F_- \frac{p_{x_0}}{2m} - \frac{\theta p_{y_0}}{\hbar} \epsilon_{\times} f(t) \cosh f(t) \\ &- \frac{\theta p_{x_0}}{\hbar} [f(t) \sinh f(t) + \epsilon_{+} f(t) \cosh f(t)] \quad (34) \end{aligned}$$

Interestingly, we find that the NC nature of spacetime affects only those particles which are in motion at time

$t = 0$. Further, the presence of $\frac{1}{\hbar}$ factor in the NC correction even after the computation of the expectation value indicates that the NC effect is inherently quantum mechanical in nature. To make the connection with the well known classical result, we first set $f(t) = f_0$ (f_0 being a small constant amplitude of GW) and $y_0 = p_{y_0} = 0$. From Eq(s) (33) and (34), we can immediately see

$$\langle x(t) \rangle = \left(x_0 + \frac{p_{x_0} t}{m} \right) + \epsilon_{+} f_0 \left(x_0 - \frac{p_{x_0} t}{m} \right) + \frac{\theta p_{x_0}}{\hbar} \epsilon_{\times} f_0 \quad (35)$$

$$\langle y(t) \rangle = \epsilon_{\times} f_0 \left(x_0 - \frac{p_{x_0} t}{m} \right) - \frac{\theta p_{x_0}}{\hbar} (\epsilon_{+} + f_0) f_0 . \quad (36)$$

Now putting $\theta = 0$ gives us the well known classical result²⁵ in the low-velocity, long-wavelength limit. From the above expressions for the expectation values, we find that the NC effect for a particle interacting with GW appears in the form of a product of θ and f_0 and hence it would be difficult to detect this effect as it is very small compared to the term showing the effect of the passing GW. This is due to the fact that noncommutativity has been introduced by Bopp-shifting the coordinates which couple with the Γ -term in the Hamiltonian and hence the NC correction in the Hamiltonian (8) contains a Γ -term. However, the present exercise gives us a substantial idea of what to expect if the same computational scheme is applied for a harmonic oscillator system. The NC effect is expected to be more prominent in this case since along with the coupling term of GW with noncommutativity, there will also be NC corrections coming from the harmonic oscillator potential which may give rise to a NC effect of comparable magnitude with GW interaction. Interaction of GW with a Hydrogen atom in the NC setting will be more involved¹⁰ and is yet another interesting case to study where a detectable NC effect

should occur. Work in this direction is in progress and will be taken up in the subsequent papers.

was completed. The authors would also like to thank the referee for useful comments.

Acknowledgment

The authors would like to acknowledge the hospitality of IIT Kanpur where a considerable part of this work

-
- * Electronic address: anirban@iucaa.ernet.in
† Electronic address: sunandan@bose.res.in, sunandan.gangopadhyay@iitk.ac.in
- ¹ A. Abramovici *et al*, Science **256** (1992) 325; <http://www.ligo.org>.
 - ² B. Caron *et al*, Class. Quant. Grav. **14** (1997) 1461; <http://www.virgo.infn.it>.
 - ³ H. Lück *et al*, Class. Quant. Grav. **14** (1997) 1471; <http://www.geo600.uni-hannover.de>.
 - ⁴ M. Ando *et al*, Phys. Rev.Lett. **86** (2001) 3950; <http://tamago.mtk.nao.ac.jp>.
 - ⁵ C. M. Caves, K. S. Thorne, R. W. P. Drever, V. D. Sandberg, M. Zimmermann, *Rev. Mod. Phys.* **52** 341 (1980).
 - ²⁵ S.M. Carroll, J.A. Harvey, V.A. Kostelecký, C.D. Lane, T. Okamoto, Phys. Rev. Lett. **87** (2001). 141601.
 - ⁷ O. Bertolami, J. G. Rosa, C. M. L. de Aragao, P. Castorina, D. Zappala, Phys. Rev. **D 72** (2005) 025010.
 - ⁸ R. Banerjee, B. Dutta Roy, S. Samanta Phys. Rev. **D 74** 045015 (2006).
 - ⁹ A. Saha, Eur. Phys. J.**C 51** 199 (2007).
 - ¹⁰ A. Stern, Phys. Rev. Lett. **100** (2008) 061601.
 - ¹¹ I. Mocioiu, M. Pospelov, and R. Roiban, Phys. Lett. B **489**, 390 (2000).
 - ¹² M. Chaichian, M.M. Sheikh-Jabbari, and A. Tureanu, Phys. Rev. Lett. **86**, 2716 (2001).
 - ¹³ See for example, R.J. Szabo, Phys. Rept. **378** (2003) 207 and the references therein.
 - ¹⁴ P. Mukherjee, A. Saha, Phys. Rev. **D 74** (2006) 027702 and the references therein.
 - ¹⁵ R.Banerjee, P. Mukherjee, S. Samanta, Phys. Rev. **D 75** :125020, 2007.
 - ¹⁶ M. M. Sheikh-Jabbari, Phys. Lett. **B 455**, 129 (1999); *Mod.Phys.Lett.* **84**, 5625 (2000).
 - ¹⁷ M. M. Najafabadi, Phys. Rev. **D 74** 025021 (2006).
 - ¹⁸ S. A. Abel, J. Jaeckel, V. V. Khoze, A. Ringwald, JHEP **0601** (2006) 105.
 - ¹⁹ B. Melic, K. Passek-Kumericki, J. Trampetic, Phys. Rev. **D 72** 057502 (2005).
 - ²⁰ C. D. Carone, J. Phys. Conf. Ser. **3796** (2006) and the references therein.
 - ²¹ P. M. Ho, H. C. Kao, Phys. Rev. Lett **88** (2002) 151602.
 - ²² A. D. Spiliotopoulos, Phys. Rev. **D 51** (1995) 1701.
 - ²³ Michele Maggiore, *Gravitational Wave, Vol I, Theory and Experiments* Oxford University Press, 2008.
 - ²⁴ <http://www.lisa-science.org>.
 - ²⁵ S. M. Carroll, *Space time Geometry: An Introduction to General Relativity*, Addison Wesley.
 - ²⁶ R. Banerjee, *Mod.Phys.Lett.*A17:631,2002.
 - ²⁷ As usual, latin indices run from 1 – 3.
 - ²⁸ Note that this condition is satisfied by resonant bar detectors and earth bound interferometers but not by the proposed space-borne interferometers such as LISA²⁴ or by the Doppler tracking of spacecraft.
 - ²⁹ We would like to mention that it is possible to shift the noncommutativity from the coordinates to the momenta leading to a dual description as shown in the literature²⁶.
 - ³⁰ x is the eigen value of the position operator X and $\vec{p}$ is the eigenvalue of the momentum operator $\vec{P}$ and $l_0 = \sqrt{\frac{\hbar}{m\omega}}$ sets the length scale.